

ICD10 codes for 27486 and 27487

| | |
|----------------------------|---|
| T84.012-13 with A/D/S | Broken |
| T84.022-23 with A/D/S | Dislocated |
| T84.032-33 with A/D/S | Loose |
| T84.042-43 with A/D/S | Periprosthetic fracture |
| T84.052-53 with A/D/S | Osteolysis |
| T84.062-63 with A/D/S | Bearing wear |
| T84.092-93 with A/D/S | Other |
| T84.099 with A/D/S | Unspecified |
| T84.53X and 54X with A/D/S | Infection |
| M89.761, M89.762, M89.769 | Major osseous defect lower leg |
| T84.7XXA (and rarely D/S) | Initial encounter for infected prosthesis (needs additional code for infecting org) |
| T84.9XX with A/D/S | Encounter for unspecified complication of prosthetic device |
| T84.10 with A/D/S | Breakdown of muscle and tendon graft (allograft ext. mechanism) failure) |
| T84.418 with A/D/S | Breakdown of implant |
| T84.420 with A/D/S | Displacement of muscle and tendon graft (allograft ext. mechanism) |
| T84.428 with A/D/S | Displacement of implant |
| T84.498 with A/D/S | Other mechanical complication of implant |
| T84.82X with A/D/S | Fibrosis due to implant |
| T84.83X with S | Recurrent hemorrhage |
| T84.84X with S | Persistent pain |
| T84.89X with S | Persistent complication |
| M61.561, M61.562, M61.569 | Other ossification of muscle, lower leg |
| M76.30, M76.31, M76.32 | Iliotibial band tendonitis (lateral impingement) |