

ICD10 codes for 27134, 27137, 27138

T84.010-11 with A/D/S	Broken
T84.020-21 with A/D/S	Dislocated
T84.030-31 with A/D/S	Loose
T84.040-41 with A/D/S	Periprosthetic fracture
T84.050-51 with A/D/S	Osteolysis
T84.060-61 with A/D/S	Bearing wear
T84.090-91 with A/D/S	Other
T84.099 with A/D/S	Unspecified
T84.50X-52X with A/D/S	Infection
T56.2X1 with A/D/S	Chromium toxicity
T56.91X with A/D/S	Cobalt toxicity
M89.751-759	Major osseous defect pelvis
T84.7XXA (and rarely D/S)	Initial encounter for infected prosthesis (needs additional code for infecting org)
T84.9XX with A/D/S	Encounter for unspecified complication of prosthetic device
S73.001-S73.046 with A/D/S	Dislocation of hip
T84.418 with A/D/S	Breakdown of implant
T84.428 with A/D/S	Displacement of implant
T84.498 with A/D/S	Other mechanical complication of implant
T84.82X with A/D/S	Fibrosis due to implant
T84.83X with S	Recurrent hemorrhage
T84.84X with S	Persistent pain
T84.89X with S	Persistent complication (Clicking, squeaking)